

## Spring 2009 CH 302:

### 30 Practice Problems Covering the Equilibrium Problem Types You Will Find on Exam 2.

- 1) The pH of an equimolar solution of acetic acid + sodium acetate is found to be 4.75.  
What will happen to the pH of this solution if we dilute it with water to twice the initial volume?
- A. The pH will rise and be higher than 4.75.
  - B. The pH will fall and be lower than 4.75.
  - C. The pH will remain 4.75.
  - D. No answer is possible without knowing the actual initial concentrations.
- 2) The pH of a solution that is 0.25 M  $(\text{NH}_4)_2\text{SO}_4(\text{aq})$  and 0.50 M  $\text{NH}_3(\text{aq})$  is:
- A. 4.75
  - B. 8.95
  - C. 9.25
  - D. 9.56
- 3) What will be the pH in a titration in which 5.0 mL of 0.120 M  $\text{HNO}_3(\text{aq})$  is added to 25.0 mL of 0.240 M  $\text{KOH}(\text{aq})$ ?
- A. 0.74
  - B. 13.26
  - C. 13.33
  - D. 13.38
- 4) The volume, in mL, calculated to one decimal point, of 0.25 M  $\text{HCl}(\text{aq})$  required to reduce the pH of 50 mL of a 0.40 M ammonia solution to a value of 7.00 would be:
- A. 80.0
  - B. 79.6
  - C. 50.0
  - D. 49.8
- 5) A buffer solution of volume 200.0 mL is 0.250 M  $\text{Na}_2\text{HPO}_4(\text{aq})$  and 0.250 M  $\text{KH}_2\text{PO}_4(\text{aq})$ . The pH resulting from the addition of 50.0 mL of 0.100 M  $\text{NaOH}(\text{aq})$  to the buffer solution will be

- A. 7.12
- B. 7.21
- C. 7.30
- D. 12.77

6) After the addition of 150 mL of 0.100 M KOH(aq) to 100 mL of 0.100 M H<sub>2</sub>SO<sub>4</sub>(aq),

- A. the pH of the final solution will be 1.92.
- B. the pH of the final solution will be greater than 1.92, but less than 7.00.
- C. the pH of the final solution will be 7.00.
- D. the pH of the final solution will be greater than 7.00, but less than 13.00.

7) The molar concentration (to two significant figures) of chloride ions in a saturated solution of silver chloride is:

- A.  $0.8 \times 10^{-10} \text{ mol}\cdot\text{L}^{-1}$
- B.  $1.6 \times 10^{-10} \text{ mol}\cdot\text{L}^{-1}$
- C.  $0.8 \times 10^{-5} \text{ mol}\cdot\text{L}^{-1}$
- D.  $1.3 \times 10^{-5} \text{ mol}\cdot\text{L}^{-1}$

8) The molar concentration (to two significant figures) of iodide ions in a saturated solution of lead iodide is:

- A.  $1.4 \times 10^{-8} \text{ mol}\cdot\text{L}^{-1}$
- B.  $1.2 \times 10^{-4} \text{ mol}\cdot\text{L}^{-1}$
- C.  $1.5 \times 10^{-3} \text{ mol}\cdot\text{L}^{-1}$
- D.  $3.0 \times 10^{-3} \text{ mol}\cdot\text{L}^{-1}$

9) What is the solubility of Ag<sub>2</sub>CrO<sub>4</sub> ( $K_{\text{sp}} = 1.12 \times 10^{-12}$ ) in grams per 100 mL of 0.100 M solution AgNO<sub>3</sub>?

- A.  $1.5 \times 10^{-9}$
- B.  $3.2 \times 10^{-8}$
- C.  $3.7 \times 10^{-9}$
- D.  $1.8 \times 10^{-8}$

- 10) When sodium nitrite is added to  $\text{HNO}_2(\text{aq})$ ,
- A. the equilibrium concentration of  $\text{HCOOH}(\text{aq})$  decreases.
  - B. the pH of the solution increases.
  - C. the  $K_a$  increases.
  - D. the pH of the solution does not change.
  - E. the pH of the solution decreases.
- 11) 100 mL of each of the following solutions is mixed; which one of the mixed solutions is a buffer?
- A. 1.0 M  $\text{NH}_3(\text{aq})$  + 0.6 M  $\text{KOH}(\text{aq})$
  - B. 1.0 M  $\text{NH}_4\text{Cl}(\text{aq})$  + 1.0 M  $\text{KOH}(\text{aq})$
  - C. 1.0 M  $\text{NH}_3(\text{aq})$  + 0.4 M  $\text{HCl}(\text{aq})$
  - D. 1.0 M  $\text{NH}_4\text{Cl}(\text{aq})$  + 0.4 M  $\text{HCl}(\text{aq})$
  - E. 1.0 M  $\text{NH}_3(\text{aq})$  + 1.0 M  $\text{HCl}(\text{aq})$
- 12) Choose the effective pH range of an aniline/anilinium chloride buffer. The value of the  $K_b$  for aniline is  $4.3 \times 10^{-10}$ .
- A. 3.6–5.6
  - B. 8.4–10.4
  - C. 1.1–3.1
  - D. 5.1–7.1
  - E. 10.1–12.1
- 13) Which of the following mixtures gives a buffer with a pH greater than 7.0? For  $\text{HCNO}$ ,  $K_a = 2.2 \times 10^{-4}$  and for  $\text{NH}_3$ ,  $K_b = 1.8 \times 10^{-5}$ .
- A. 10 mL of 0.1 M  $\text{NH}_3(\text{aq})$  + 10 mL of 0.1 M  $\text{HCl}(\text{aq})$
  - B. 10 mL of 0.1 M  $\text{HCNO}(\text{aq})$  + 10 mL of 0.1 M  $\text{NaOH}(\text{aq})$
  - C. 10 mL of 0.1 M  $\text{HCNO}(\text{aq})$  + 5.0 mL of 0.1 M  $\text{NaOH}(\text{aq})$
  - D. 10 mL of 0.1 M  $\text{NH}_3(\text{aq})$  + 10 mL of 0.1 M  $\text{HCNO}(\text{aq})$
  - E. 10 mL of 0.1 M  $\text{NH}_3(\text{aq})$  + 5.0 mL of 0.1 M  $\text{HCl}(\text{aq})$

14) If a small amount of a strong base is added to buffer made up of a weak acid, HA, and the sodium salt of its

conjugate base, NaA, the pH of the buffer solution does not change appreciably because

- A. the  $K_a$  of HA is changed.
- B. No reaction occurs.
- C. the strong base reacts with  $A^-$  to give HA, which is a weak acid.
- D. the strong base reacts with HA to give AOH and  $H^+$ .
- E. the strong base reacts with HA to give  $A^-$ , which is a weak base.


15) At the stoichiometric point in the titration of 0.130 M HCOOH(aq) with 0.130 M KOH(aq),

- A. the pH is 7.0.
- B.  $[HCOOH] = 0.0650$  M.
- C.  $[HCO_2^-] = 0.130$  M.
- D. the pH is greater than 7.
- E. the pH is less than 7.

16) For the titration of 50.0 mL of 0.020 M aqueous salicylic acid with 0.020 M KOH(aq), calculate the pH after the addition of 55.0 mL of KOH(aq). For salicylic acid,  $pK_a = 2.97$ .

- A. 10.98
- B. 7.00
- C. 11.26
- D. 12.02
- E. 12.30


17) The titration curve for the titration of 0.100 M  $H_2SO_3$ (aq) with 0.100 M KOH(aq) is given below.



Estimate  $pK_{a1}$  and  $pK_{a2}$  of  $H_2SO_3$ .

Answer

18) The titration curve for the titration of 0.100 M  $\text{Na}_2\text{CO}_3(\text{aq})$  with 0.100 M  $\text{HClO}_4(\text{aq})$  is:



The main species in the solution after the addition of 35 mL of  $\text{HClO}_4$  are

- A.  $\text{HCO}_3^-$ ,  $\text{H}_2\text{CO}_3$ ,  $\text{Na}^+$ , and  $\text{ClO}_4^-$ .
- B.  $\text{H}_2\text{CO}_3$ ,  $\text{Na}^+$ , and  $\text{ClO}_4^-$ .
- C.  $\text{CO}_3^{2-}$ ,  $\text{HCO}_3^-$ ,  $\text{Na}^+$ , and  $\text{ClO}_4^-$ .
- D.  $\text{CO}_3^{2-}$ ,  $\text{Na}^+$ , and  $\text{ClO}_4^-$ .
- E.  $\text{HCO}_3^-$ ,  $\text{Na}^+$ , and  $\text{ClO}_4^-$ .

19) What is the relationship between the solubility in water,  $s$ , and  $K_{\text{sp}}$  for the ionic solid  $\text{Ca}_3(\text{PO}_4)_2$ ?

- A.  $K_{\text{sp}} = 72s^5$
- B.  $K_{\text{sp}} = 5s$
- C.  $K_{\text{sp}} = 6s^2$
- D.  $K_{\text{sp}} = s^5$

20) Which of the following water-insoluble salts is more soluble in 1.0 M  $\text{HClO}_4(\text{aq})$ ?

- A.  $\text{AgBr}$
- B.  $\text{PbF}_2$
- C.  $\text{Hg}_2\text{Br}_2$
- D.  $\text{PbI}_2$
- E.  $\text{AgClO}_4$

21) If equal volumes of 0.004 M  $\text{Pb}(\text{NO}_3)_2(\text{aq})$  and 0.004 M  $\text{KI}(\text{aq})$  are mixed, what reaction, if any, occurs? The value of  $K_{\text{sp}}$  for  $\text{PbI}_2$  is  $1.4 \times 10^{-8}$ .

- A. The solution turns purple because of formation of  $\text{I}_2$ .
- B.  $\text{PbI}_2(\text{s})$  precipitates.
- C.  $\text{KNO}_3(\text{s})$  precipitates.
- D. No reaction occurs.
- E. The value of  $K_{\text{sp}}$  changes to  $9 \times 10^{-9}$ .

22) If you wish to increase the solubility of silver benzoate, a preservative, you would

- A. add sodium hydroxide.
- B. decrease the pH.
- C. add sodium acetate.
- D. add sodium benzoate.
- E. add silver nitrate.

23) Rank, from greater to least, the equilibrium concentration of species formed when 0.2M  $\text{H}_2\text{SO}_4$  is placed in solution.

Answer

24) You have watched a fellow student derive the exact solution for a dilute strong acid in water. Feel free to derive a similar solution for a dilute weak acid case (or look at the equation in the notes.) How many more unknowns are there for the dilute weak acid than the dilute weak acid? What is the number of coefficients in the charge balance equation for weak acid case? How

Answer

25) Consider the following solutions made by dumping a polyprotic acid,  $\text{H}_3\text{Y}$ , or its salts into water (hint, get rid of those pesky spectators):

Solution I:  $\text{H}_3\text{Y}$

Solution II:  $\text{Li}_2\text{NaY}$

Solution III:  $\text{K}_2\text{HY}$  and  $\text{K}_2\text{NaY}$

Solution IV:  $\text{CaHY}$

What are the simple acid/base equations you would use to find the pH for these solutions?

Answer

26) A solution is made by placing 100 ml of 0.3M ammonia and 200 ml of 0.5M ammonium nitrate in water. What is the buffer capacity for this solution?

Answer

27) Place a drop of aqueous HCl on chalk (calcium carbonate) and watch the fizzing begin. Name all the chemical species formed from the carbonate and their associated chemical equilibria.

Answer

28) Consider a solution made by placing  $10^{-8}$  M KOH and 0.1 M ammonia in water. Rank from most to least, the species that provide the greatest number of hydroxides.

- A. water > KOH > ammonia
- B. KOH > ammonia > water
- C. KOH > water > ammonia
- D. ammonia ? water > KOH

29) Assuming equal amounts of each are placed in water, which of the following compounds produces the strongest acid?

- A. The acid of sodium formate which has a  $K_b$  of  $10^{-9}$ ?
- B. Acetic acid which has a salt with a  $K_b$  of  $5 \times 10^{-10}$ ?
- C. Pure water
- D. Tartaric acid which has a  $K_a$  of  $1 \times 10^{-6}$
- E. Ammonia which has a salt with a  $K_a$  of  $5 \times 10^{-10}$ ?

30) Which of the following solutions would not require more than one equilibrium constant to accurately calculate pH?

- A. A mixture of methanol and water.
- B. A 0.1 M solution of sulfuric acid.
- C. A solution of sodium bicarbonate.
- D. A  $10^{-6}$  M solution of ammonia?
- E. An equimolar solution of  $\text{Na}_2\text{LiPO}_4$  and  $\text{Na}_2\text{HPO}_4$ ?